



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

NOTE ON A SUBSTITUTE FOR DUHAMEL'S THEOREM.

BY HENRY B. FINE.

The substitute for Duhamel's Theorem published by G. A. Bliss in the sixteenth volume of this journal* meets all the requirements of the case for students who are mature enough mathematically to understand it. But it is cast in too abstract a form to be serviceable as early in the study of the calculus as some theorem of this kind must be used. A less general theorem is all that is needed, but both in statement and proof it must be sufficiently elementary to be intelligible to a student soon after he has become familiar with the notion of the simple definite integral as the limit of a sum of infinitesimal elements and has acquired some skill in applying this notion. I am therefore led to suggest the following theorem. In content it is the equivalent of a theorem recently published by E. V. Huntington† and like that is a particular case of Bliss's theorem, but the proof, unlike Huntington's and like Bliss's, shows directly the existence of the limit of the sum in question.

THEOREM. *Let $f_1(x), f_2(x), \dots, f_p(x)$ denote any set of functions of x , finite in number, which are continuous in the interval (ab) , and let*

$$F(x) = f_1(x)f_2(x) \cdots f_p(x).$$

Suppose the interval (ab) to be divided and redivided into parts in any manner such that as the process is indefinitely continued the greatest of the parts will approach 0 as limit, and at any stage in the process let h_1, h_2, \dots, h_n represent the parts in length and position; also let $\xi'_1, \xi''_1, \dots, \xi^{(p)}_1$, and ξ_i denote any numbers in the part h_i ($i = 1, 2, \dots, n$). Then

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n f_1(\xi'_i)f_2(\xi''_i) \cdots f_p(\xi^{(p)}_i)h_i = \lim_{n \rightarrow \infty} \sum_{i=1}^n F(\xi_i)h_i = \int_a^b F(x)dx.$$

For any difference of the form $a_1a_2 \cdots a_p - b_1b_2 \cdots b_p$ can be expressed in terms of the differences $a_i - b_i$ by means of the identity

$$\begin{aligned} a_1a_2 \cdots a_p - b_1b_2 \cdots b_p &= (a_1 - b_1)a_2a_3 \cdots a_p + (a_2 - b_2)b_1a_3 \cdots a_p \\ &\quad + (a_3 - b_3)b_1b_2a_4 \cdots a_p + \cdots + (a_p - b_p)b_1b_2 \cdots b_{p-1}. \end{aligned}$$

* G. A. Bliss, "A Substitute for Duhamel's Theorem," *Annals of Mathematics*, Ser. 2, vol. 16 (1914).

† E. V. Huntington, "On Setting Up a Definite Integral without the Use of Duhamel's Theorem," *The American Mathematical Monthly*, vol. 24 (1917).

Applying this identity to the difference

$$f_1(\xi_i')f_2(\xi_i'') \cdots f_p(\xi_i^{(p)}) - f_1(\xi_i)f_2(\xi_i) \cdots f_p(\xi_i)$$

and in the coefficient of each difference $f_j(\xi_i^{(j)}) - f_j(\xi_i)$ on the right replacing every factor f by M , the greatest of the maximum absolute values of the several functions $f_j(x)$ in (ab) , we have

$$\begin{aligned} |f_1(\xi_i')f_2(\xi_i'') \cdots f_p(\xi_i^{(p)}) - F(\xi_i)| &\leq |f_1(\xi_i') - f_1(\xi_i)| M^{p-1} \\ &+ |f_2(\xi_i'') - f_2(\xi_i)| M^{p-1} + \cdots + |f_p(\xi_i^{(p)}) - f_p(\xi_i)| M^{p-1}. \end{aligned}$$

But since all the functions $f_j(x)$ are continuous, and therefore uniformly continuous, in (ab) , if any positive number ϵ be assigned we can find an integer n' such that for $n > n'$, and $i = 1, 2, \dots, n$, we shall have

$$\begin{aligned} |f_1(\xi_i') - f_1(\xi_i)| &< \frac{\epsilon}{pM^{p-1}}, \quad |f_2(\xi_i'') - f_2(\xi_i)| < \frac{\epsilon}{pM^{p-1}}, \quad \dots, \\ |f_p(\xi_i^{(p)}) - f_p(\xi_i)| &< \frac{\epsilon}{pM^{p-1}}, \end{aligned}$$

and therefore, by the above inequality,

$$|f_1(\xi_i')f_2(\xi_i'') \cdots f_p(\xi_i^{(p)}) - F(\xi_i)| < \epsilon \quad (i = 1, 2, \dots, n).$$

From this it immediately follows, since

$$\sum_{i=1}^n h_i = b - a,$$

that

$$\left| \sum_{i=1}^n f_1(\xi_i')f_2(\xi_i'') \cdots f_p(\xi_i^{(p)})h_i - \sum_{i=1}^n F(\xi_i)h_i \right| < \epsilon(b - a),$$

and therefore, since $\epsilon(b - a)$ may be taken as small as we please, that

$$\lim_{n \rightarrow \infty} \left\{ \sum_{i=1}^n f_1(\xi_i')f_2(\xi_i'') \cdots f_p(\xi_i^{(p)})h_i - \sum_{i=1}^n F(\xi_i)h_i \right\} = 0.$$

Hence since $\lim_{n \rightarrow \infty} \sum_{i=1}^n F(\xi_i)h_i$ exists, and by definition is $\int_a^b F(x)dx$, the

theorem is proved.

The theorem and its proof can be immediately extended to the more general integral

$$\int_V F(x_1, x_2, \dots, x_m) dV$$

where F denotes a product of p functions f_j all of which are supposed continuous in the closed region V . It is only necessary in the proof just given to replace h_i by the element ΔV_i of V and to interpret the several ξ_i 's as symbols for sets of values of x_1, x_2, \dots, x_m in ΔV_i .